Author index

Abiko, Y., see Sugiyama, N. (952) 1
Alemany, S., see Cabrero, C. (952) 277
Anderson, V.E., see Wlassics, I.D. (952) 269
Anthony, F.A., see Fong, K.C. (952) 13
Arrondo, J.L.R., Young, N.M. and Mantsch, H.H.
The solution structure of concanavalin A probed by FT-IR spectroscopy (952) 261
Asakura, S., Yoshida, N., Matsuda, M., Murayama, H. and Soe, G.
Preparation and characterization of monoclonal antibodies against the human thrombin-antithrombin III complex (952) 37
Autuori, F., see Beninati, S. (952) 325
Avizonis, D.Z., see Satterlee, J.D. (952) 317

Babitch, J.A., see Fong, K.C. (952) 13
Bardakdjian-Michau, J., see Rhoda, M.-D. (952) 208
Beninati, S., Piacentini, M., Cocuzzi, E.T., Autuori, F. and Folk, J.E.
Covalent incorporation of polyamines as γ-glutamyl derivatives into CHO cell protein (952) 325
Beuzard, Y., see Rhoda, M.-D. (952) 208
Burton, J. and Quinn, T.
The amino-acid residues on the C-terminal side of the cleavage site of angiotensinogen influence the species specificity of reaction with renin (952) 8

Cabrero, C. and Alemany, S.
Conversion of rat liver *S*-adenosyl-L-methionine synthetase from high-M_r form to low-M_r form by LiBr (952) 277
Caughey, G.H., Viro, N.F., Lazarus, S.C. and Nadel, J.A.
Purification and characterization of dog mastocytoma chymase: identification of an octapeptide conserved in chymotryptic leukocyte proteinases (952) 142
Citri, N., see Samuelov, N.S. (952) 48
Cocuzzi, E.T., see Beninati, S. (952) 325
Cole, T.-C., Melrose, J. and Ghosh, P.
Cysteine proteinase inhibitors of the canine intervertebral disc (952) 201
Cramer, S.P., see Prince, R.C. (952) 220

De Arriaga, D., see Montero, S. (952) 56
De Silva, D.G.A.H. and Sykes, A.G.
Interpretation of effects of pH on rate constants for the oxidation of three ferrocytochromes *c*-551 with $[Fe(CN)_6]^{3-}$ and $[Co(phen)_3]^{3+}$, and assignment of pK_a values (952) 334
Dickson, D.P.E., see St. Pierre, T.G. (952) 158
Domenget, C., see Rhoda, M.-D. (952) 208

Eady, R.R., see Miller, R.W. (952) 290
Eberspach, I., Straßburger, W., Glatter, U., Gerday, C. and Wollmer, A.
Interaction of parvalbumin of pike II with calcium and terbium ions (952) 67
Endo, A., see Sakai, K. (952) 297
Epand, R.M., see Kahan, I. (952) 230

Fay, P.J.
Subunit structure of thrombin-activated human Factor VIIIa (952) 181
Folk, J.E., see Beninati, S. (952) 325
Fong, K.C., Babitch, J.A. and Anthony, F.A.
Calcium binding to tubulin (952) 13
Foster, D., see Petersen, L.C. (952) 245
Frömmel, C., see Hausdorf, G. (952) 20
Fukasawa, S., see Yamakura, F. (952) 304
Fukui, T., Narikawa, T., Miwa, K., Shirakura, Y., Saito, T. and Tomita, K.
Effect of limited tryptic modification of a bacterial poly(3-hydroxybutyrate) depolymerase on its catalytic activity (952) 164
Funasaki, N., see Neya, S. (952) 150

George, G.N., see Prince, R.C. (952) 220
Gerday, C., see Eberspach, I. (952) 67
Ghosh, P., see Cole, T.-C. (952) 201
Glatter, U., see Eberspach, I. (952) 67

Hägeli, G., see Pellegrini, A. (952) 309
Hasumi, K., see Sakai, K. (952) 297
Hausdorf, G., Krüger, K., Küttner, G., Holzhütter, H.-G., Frömmel, C. and Höhne, W.E.
Oxidation of a methionine residue in subtilisin-type proteinases by the hydrogen peroxide/borate system — an active site-directed reaction (952) 20
Hawkes, J.J. and Pethig, R.
Low-frequency dielectric properties of lysozyme as a function of hydration and pH of lyophilisation (952) 27
Hoggett, J.G., see Woolfitt, A.R. (952) 238
Höhne, W.E., see Hausdorf, G. (952) 20
Holzhütter, H.-G., see Hausdorf, G. (952) 20

Ichikawa, M. and Toraya, T.
Roles of the β-D-ribofuranose ring and the functional groups of the D-ribose moiety of adenosylcobalamin in the diol dehydratase reaction (952) 191
Ito, S., see Shikata, S. (952) 282
Iwamoto, M., see Sugiyama, N. (952) 1

Johannessen, M., see Petersen, L.C. (952) 245
Jue, T., see La Mar, G.N. (952) 131

Kahan, I., Epand, R.M. and Moscarello, M.A.
The secondary structure of a membrane-embedded peptide from the carboxy terminus of lipophilin as revealed by circular dichroism (952) 230
Kauten, R.J., see La Mar, G.N. (952) 131
Kawai, S., see Shikata, S. (952) 282
Kellett, G.L., see Woolfitt, A.R. (952) 238
Knull, H.R., see Walsh, J.L. (952) 83
Kobayashi, H., see Yamakura, F. (952) 304
Krenn, B.E., Plat, H. and Wever, R.
Purification and some characteristics of a non-haem bromoperoxidase from *Streptomyces aureofaciens* (952) 255
Krüger, K., see Hausdorf, G. (952) 20
Kumar, A., see Petersen, L.C. (952) 245
Küttner, G., see Hausdorf, G. (952) 20

La Mar, G.N., Jue, T., Nagai, K., Smith, K.M., Yamamoto, Y., Kauten, R.J., Thanabal, V., Langry, K.C., Pandey, R.K. and Leung, H.-K.
^{1}H-NMR heme resonance assignments by selective deuteration in low-spin complexes of ferric hemoglobin A (952) 131
Langry, K.C., see La Mar, G.N. (952) 131
Lazarus, S.C., see Caughey, G.H. (952) 142
Leung, H.-K., see La Mar, G.N. (952) 131

Mäenpää, P.H., see Palvimo, J. (952) 172
Mantsch, H.H., see Arrondo, J.L.R. (952) 261
Mantsch, H.H., see Surewicz, W.K. (952) 115
Marchese-Ragona, S.P., see Merkler, D.J. (952) 101
Matsuda, M., see Asakura, S. (952) 37
Melrose, J., see Cole, T.-C. (952) 201
Merkler, D.J., Srikumar, K., Marchese-Ragona, S.P. and Wedler, F.C.
Aggregation and thermo-inactivation of glutamine synthetase from an extreme thermophile, *Bacillus caldolyticus* (952) 101
Miller, R.W. and Eady, R.R.
Cyanamide: a new substrate for nitrogenase (952) 290
Miwa, K., see Fukui, T. (952) 164
Moench, S.J., see Satterlee, J.D. (952) 317
Montero, S., De Arriaga, D. and Soler, J.
Purification and characterization of glutathione reductase from mycelium of *Phycomyces blakesleeanus* (952) 56
Moscarello, M.A., see Kahan, I. (952) 230
Mulvihill, E., see Petersen, L.C. (952) 245
Murayama, H., see Asakura, S. (952) 37

Nadel, J.A., see Caughey, G.H. (952) 142
Nagai, K., see La Mar, G.N. (952) 131
Narikawa, T., see Fukui, T. (952) 164
Neya, S. and Funasaki, N.
Proton NMR study of the myoglobin reconstituted with *meso*-tetra(*n*-propyl)hemin (952) 150

Okamoto, K., see Shikata, S. (952) 282
Ozaki, K., see Shikata, S. (952) 282

Palvimo, J. and Mäenpää, P.H.
Binding of high-mobility-group proteins HMG 14 and HMG 17 to DNA and histone H1 as influenced by phosphorylation (952) 172
Pandey, R.K., see La Mar, G.N. (952) 131
Patel, R.N., see Prince, R.C. (952) 220
Pellegrini, A., Hägeli, G. and Von Fellenberg, R.
Isolation and characterization of two new low-molecular-weight protein proteinase inhibitors from the granule-rich fraction of equine neutrophilic granulocytes (952) 309
Peters, T.J., see St. Pierre, T.G. (952) 158
Petersen, L.C., Johannessen, M., Foster, D., Kumar, A. and Mulvihill, E.
The effect of polymerised fibrin on the catalytic activities of one-chain tissue-type plasminogen activator as revealed by an analogue resistant to plasmin cleavage (952) 245
Pethig, R., see Hawkes, J.J. (952) 27
Piacentini, M., see Beninati, S. (952) 325
Pick, E., see Sha'ag, D. (952) 213
Plat, H., see Krenn, B.E. (952) 255
Pollard, R.K., see St. Pierre, T.G. (952) 158
Prince, R.C., George, G.N., Savas, J.C., Cramer, S.P. and Patel, R.N.
Spectroscopic properties of the hydroxylase of methane monooxygenase (952) 220

Quinn, T., see Burton, J. (952) 8

Rhoda, M.-D., Domenget, C., Vidaud, M., Bardakdjian-Michau, J., Rouyer-Fessard, P., Rosa, J. and Beuzard, Y.
Mouse α chains inhibit polymerization of hemoglobin induced by human β^S or $\beta^{S\,Antilles}$ chains (952) 208
Rigo, A., see Viglino, P. (952) 77
Rosa, J., see Rhoda, M.-D. (952) 208
Rotilio, G., see Viglino, P. (952) 77
Rouyer-Fessard, P., see Rhoda, M.-D. (952) 208

Saito, T., see Fukui, T. (952) 164
Sakai, K., Hasumi, K. and Endo, A.
Inactivation of rabbit muscle glyceraldehyde-3-phosphate dehydrogenase by koningic acid (952) 297
Samuelov, N.S. and Citri, N.
Inducible oxacillin-hydrolyzing β-lactamase in a methylotrophic bacterium (952) 48
Sasaki, T., see Sugiyama, N. (952) 1
Satterlee, J.D., Avizonis, D.Z. and Moench, S.J.
Assignment of hyperfine-shifted resonances in yeast ferricytochrome *c* isozyme 2 using the proton pre-steady-state nuclear Overhauser effect (952) 317
Savas, J.C., see Prince, R.C. (952) 220
Scarpa, M., see Viglino, P. (952) 77
Seehafer, J.G., Slupsky, J.R., Tang, S.-C. and Shaw, A.R.E.
The functional cell surface glycoprotein CD9 is distinguished by being the major fatty acid acylated and a major iodinated cell-surface component of the human platelet (952) 92

Sha'ag, D. and Pick, E.
Macrophage-derived superoxide-generating NADPH oxidase in an amphiphile-activated, cell-free system; partial purification of the cytosolic component and evidence that it may contain the NADPH binding site (952) 213
Shaw, A.R.E., see Seehafer, J.G. (952) 92
Shikata, S., Ozaki, K., Kawai, S., Ito, S. and Okamoto, K.
Purification and characterization of $NADP^+$-linked isocitrate dehydrogenase from an alkalophilic *Bacillus* (952) 282
Shirakura, Y., see Fukui, T. (952) 164
Slupsky, J.R., see Seehafer, J.G. (952) 92
Smith, K.M., see La Mar, G.N. (952) 131
Soe, G., see Asakura, S. (952) 37
Soler, J., see Montero, S. (952) 56
Srikumar, K., see Merkler, D.J. (952) 101
Stille, C., see Wlassics, I.D. (952) 269
St. Pierre, T.G., Pollard, R.K., Dickson, D.P.E., Ward, R.J. and Peters, T.J.
Mössbauer spectroscopic studies of deproteinised, subfractionated and reconstituted ferritins: the relationship between haemosiderin and ferritin (952) 158
Straßburger, W., see Eberspach, I. (952) 67
Sugiyama, N., Sasaki, T., Iwamoto, M. and Abiko, Y.
Binding site of α_2-plasmin inhibitor to plasminogen (952) 1
Surewicz, W.K. and Mantsch, H.H.
New insight into protein secondary structure from resolution-enhanced infrared spectra (952) 115
Sykes, A.G., see De Silva, D.G.A.H. (952) 334

Tang, S.-C., see Seehafer, J.G. (952) 92
Thanabal, V., see La Mar, G.N. (952) 131
Tomita, K., see Fukui, T. (952) 164
Toraya, T., see Ichikawa, M. (952) 191

Vidaud, M., see Rhoda, M.-D. (952) 208
Viglino, P., Scarpa, M., Rotilio, G. and Rigo, A.
A kinetic study of the reactions between H_2O_2 and Cu,Zn superoxide dismutase; evidence for an electrostatic control of the reaction rate (952) 77
Viro, N.F., see Caughey, G.H. (952) 142
Von Fellenberg, R., see Pellegrini, A. (952) 309

Walsh, J.L. and Knull, H.R.
Heteromerous interactions among glycolytic enzymes and of glycolytic enzymes with F-actin: effects of poly(ethylene glycol) (952) 83
Ward, R.J., see St. Pierre, T.G. (952) 158
Watanabe, K., see Yamakura, F. (952) 304
Wedler, F.C., see Merkler, D.J. (952) 101
Wever, R., see Krenn, B.E. (952) 255
Wlassics, I.D., Stille, C. and Anderson, V.E.
Coenzyme A dithioesters: synthesis, characterization and reaction with citrate synthase and acetyl-CoA : choline *O*-acetyltransferase (952) 269
Wollmer, A., see Eberspach, I. (952) 67
Woolfitt, A.R., Kellett, G.L. and Hoggett, J.G.
The binding of glucose and nucleotides to hexokinase from *Saccharomyces cerevisiae* (952) 238

Yamakura, F., Watanabe, K., Kobayashi, H. and Fukasawa, S.
A comparison of the bacteriocupreins in crude extracts of *Photobacterium leiognathi* isolated from squid and fish (952) 304
Yamamoto, Y., see La Mar, G.N. (952) 131
Yoshida, N., see Asakura, S. (952) 37
Young, N.M., see Arrondo, J.L.R. (952) 261